

ISSN: 1401-5617



Intersections in genus 3 and the Boussinesq hierarchy

S. V. Shadrin

RESEARCH REPORTS IN MATHEMATICS
NUMBER 11, 2003

DEPARTMENT OF MATHEMATICS
STOCKHOLM UNIVERSITY

Electronic versions of this document are available at
<http://www.math.su.se/reports/2003/11>

Date of publication: October 20, 2003

2000 Mathematics Subject Classification: Primary 14H10, Secondary 14H70, 14H81.

Keywords: Witten conjecture, Boussinesq hierarchy.

Postal address:

Department of Mathematics

Stockholm University

S-106 91 Stockholm

Sweden

Electronic addresses:

<http://www.math.su.se/>

info@math.su.se

INTERSECTIONS IN GENUS 3 AND THE BOUSSINESQ HIERARCHY

S. V. SHADRIN

ABSTRACT. In this paper we prove that the enlarged Witten's conjecture is true in the case of the Boussinesq hierarchy for correlators $\langle \tau_{n,m} \tau_{0,1}^k \tau_{0,0}^l \rangle_3$.

1. INTRODUCTION

1.1. In this paper we study a very special case of the enlarged Witten's conjecture. In [9, 10], Witten conjectured that the generating functions of certain intersection numbers $\langle \tau_{n_1, m_1} \dots \tau_{n_s, m_s} \rangle_g$ on the moduli spaces of curves are the string solutions to the n -KdV hierarchies.

This conjecture was completely proved in the case of 2-KdV hierarchy in [4] and [6] (see also [5] for some comments). For the n -KdV, $n = 3, 4, \dots$, this conjecture was proved only in genus zero ($g = 0$), see [10].

As for the higher genera, the situation looks like follows. In genera 1 and 2, there exist topological recursion relations (see [1]) allowing to calculate intersection numbers participating in the Witten's conjecture (that's why we consider genus 3 in this paper). For an arbitrary genus there exist an algorithm (see [8]) allowing to calculate (in a sense) only the intersection numbers like $\langle \tau_{n,m} \tau_{0,m_1} \dots \tau_{0,m_s} \rangle_g$ (with descendants only at one point).

1.2. In this paper we restrict ourselves to the case of 3-KdV hierarchy. We consider intersections in genus 3 with descendants only at one point. For these numbers we give some relations slightly generalizing relations from [8]. Using these relations we prove an expression of intersection numbers in genus 3 via intersection numbers in genus 0. The same expression is predicted by the 3-KdV (Boussinesq) hierarchy. Since the conjecture is already proved in genus zero, our theorem (Theorem 3) follows that the conjecture is true for the intersection numbers $\langle \tau_{n,m} \tau_{0,1}^k \tau_{0,0}^l \rangle_3$ (of course, in the case of the 3-KdV hierarchy).

The paper is organized as follows. In section 2 we recall the Witten's conjecture (the particular case we need here). In section 3 we define some auxiliary intersection numbers and give some relations. In section 4 we outline the calculation leading to the proof of the Witten's conjecture in our particular case.

Partially supported by the grants RFBR 01-01-00660 and RFBR 02-01-22004.

1.3. At the end, we have to acknowledge that we don't understand completely the original formulation of the Witten's conjecture [10] and the more rigorous formulation in [3] as well. But for our arguments it is enough to know the factorization property of the Witten's top Chern class (see definitions below) and that everything is already proved in genus zero.

This paper could be consider as a sequel to [8]. Though we have tried to make this paper be self-closed, it could be useful to look through [8] to learn some geometrical ideas hidden behind our calculations.

2. THE WITTEN'S CONJECTURE

2.1. In this section we briefly formulate the Witten's conjecture. Note that we consider here only the special case of the conjecture related to the Boussinesq hierarchy. Note also that we doesn't follow the initial Witten's definitions, but we use an axiomatic approach from [3].

2.2. Consider the moduli space $\overline{\mathcal{M}}_{g,s} \ni (C, x_1, \dots, x_s)$. Label each marked point x_i by the number $m_i \in \{0, 1\}$.

By K denote the canonical line bundle of C . Consider the line bundle $S = K \otimes (\otimes_{i=1}^s \mathcal{O}(x_i)^{-m_i})$ over C . If $2g - 2 - \sum_{i=1}^s m_i$ is divisible by 3, then there are 3^{2g} isomorphism classes of line bundles \mathcal{T} such that $\mathcal{T}^{\otimes 3} \cong S$.

The choice of an isomorphism class of \mathcal{T} determines a cover $\mathcal{M}'_{g,s}$ of $\mathcal{M}_{g,s}$. To extend it to a covering of $\overline{\mathcal{M}}_{g,s}$ we have to discuss the behavior of \mathcal{T} near a double point.

2.3. There are 3 possible cases of behavior of \mathcal{T} near the double point. Two cases we call "good", and the third case is "bad". For our purposes we need a description only of the good cases.

Let C be a singular curve with one double point. By $\pi: C_0 \rightarrow C$ denote its normalization. The preimage of the double point consists of two points, say x' and x'' . Two good cases look like follows: $\mathcal{T} \cong \pi_* \mathcal{T}'$, where \mathcal{T}' is a locally free sheaf on C_0 with a natural isomorphism either $\mathcal{T}'^{\otimes r} \cong K \otimes (\otimes_{i=1}^s \mathcal{O}(x_i)^{-m_i}) \otimes \mathcal{O}(x')^{-1}$, or $\mathcal{T}'^{\otimes r} \cong K \otimes (\otimes_{i=1}^s \mathcal{O}(x_i)^{-m_i}) \otimes \mathcal{O}(x'')^{-1}$.

2.3.1. Now we are ready to define the Witten's top Chern class $c_D(\mathcal{V}) \in H^{2D}(\overline{\mathcal{M}}'_{g,s})$, which plays the main role in the Witten's conjecture. The number D is equal to $(g - 1)/3 + \sum_{i=1}^s m_i/3$.

If $g = 0$, then $c_D(\mathcal{V})$ is the top Chern class of the bundle over $\overline{\mathcal{M}}'_{g,s}$, which fiber at a moduli point $(C, x_1, \dots, x_s, \mathcal{T})$ is equal to $H^1(C, \mathcal{T})$. In higher genera the Witten's top Chern class is just a cohomology class satisfying the following properties.

First, consider a component of the boundary of $\overline{\mathcal{M}}'_{g,s}$ consisting of curves with one double point, where \mathcal{T} is defined by a "good" case. If this component of the boundary consists of two-component curves,

then $c_D(\mathcal{V}) = c_{D_1}(\mathcal{V}_1) \cdot c_{D_2}(\mathcal{V}_2)$, where $c_{D_i}(\mathcal{V}_i)$ are the Witten's top Chern classes on the corresponding moduli spaces of the components of normalizations. If this component of the boundary consists of one-component self-intersecting curves, then $c_D(\mathcal{V}) = c_D(\mathcal{V}')$, where $c_D(\mathcal{V}')$ is defined on the normalization of these curves.

Second, consider a component of the boundary consisting of curves with one double point, where we have the "bad" case of the behavior of \mathcal{T} . In this case we require that the restriction of $c_D(\mathcal{V})$ to this component of the boundary vanishes.

2.4. These properties uniquely determine the intersection numbers, which we consider in this paper. As far as we understand, the existence of the Witten's top Chern class is proved in [7].

2.5. Let us label each marked point x_i by an integer $n_i \geq 0$. By $\langle \tau_{n_1, m_1} \dots \tau_{n_s, m_s} \rangle_g$ denote the intersection number

$$(1) \quad \langle \tau_{n_1, m_1} \dots \tau_{n_s, m_s} \rangle_g := \frac{1}{3g} \int_{\overline{\mathcal{M}}'_{g,s}} \prod_{i=1}^s \psi_i^{n_i} \cdot c_D(\mathcal{V}).$$

Here ψ_i is the first Chern class of the line bundle over $\overline{\mathcal{M}}'_{g,s}$, whose fiber at a moduli point $(C, x_1, \dots, x_s, \mathcal{T})$ is equal to $T_{x_i}^* C$. Of course, the number $\langle \tau_{n_1, m_1} \dots \tau_{n_s, m_s} \rangle_g$ is not zero only if $3g - 3 + s = \sum_{i=1}^s n_i + D$.

2.6. The special case of the conjecture from [10] is that the generating function for the numbers $\langle \tau_{n_1, m_1} \dots \tau_{n_s, m_s} \rangle_g$ is the string solution to the Boussinesq hierarchy.

In particular, for the numbers $\langle \tau_{n,m} \tau_{0,1}^k \tau_{0,0}^l \rangle_3$ the conjecture means that

$$(2) \quad \langle \tau_{n,m} \tau_{0,1}^k \tau_{0,0}^l \rangle_3 = \frac{1}{3!12^3} \langle \tau_{n-6,m} \tau_{0,1}^{k+3} \tau_{0,0}^l \rangle_0.$$

(e.g., see relation for the Boussinesq hierarchy in [8])

This is just what we prove in this paper.

3. RELATIONS FOR INTERSECTION NUMBERS

3.1. First of all, let us define some natural subvarieties of the moduli spaces of curves.

Consider the moduli space $\overline{\mathcal{M}}'_{g,s}$. We assume that the covering $\pi: \overline{\mathcal{M}}'_{g,s} \rightarrow \overline{\mathcal{M}}_{g,s}$ is defined by the collections of numbers m_1, \dots, m_s . Of course, m_i corresponds to the i -th marked point, and each m_i equals either to 0 or to 1.

Let $t < s$ be a positive integer. Consider a collection of positive integers a_1, \dots, a_t . By $F(a_1, \dots, a_t)$ denote the subvariety of the open moduli space $\mathcal{M}_{g,s}$ consisting of curves (C, x_1, \dots, x_s) such that $(-\sum_{i=1}^t a_i x_i + a_1 x_2 + \dots + a_t x_{t+1})$ is a divisor of a meromorphic function. By $F'(a_1, \dots, a_t)$ denote the closure of $\pi^{-1}(F(a_1, \dots, a_t))$ in $\overline{\mathcal{M}}'_{g,s}$.

3.2. Now let us introduce notations for the integrals over $F'(a_1, \dots, a_t)$.

Consider the numbers m_{t+2}, \dots, m_s . Let q of these numbers be equal to 0, and p of these numbers be equal to 1. Of course, $p + q = s - t - 1$. By $U(g, n, m_1, p, q | \prod_{i=1}^t \eta_{m_{i+1}, a_i})$ denote the integral of $\psi_1^n \cdot c_D(\mathcal{V})$ over the subspace $F'(a_1, \dots, a_t)$:

$$(3) \quad U(g, n, m_1, p, q | \prod_{i=1}^t \eta_{m_{i+1}, a_i}) := \frac{1}{3^g} \int_{F'(a_1, \dots, a_t)} \psi_1^n \cdot c_D(\mathcal{V}).$$

3.3. These numbers play the main role in our calculations below. In particular, there is an expression of $\langle \tau_{n,m} \tau_{0,1}^k \tau_{0,0}^l \rangle_3$ via such numbers.

Theorem 1.

$$(4) \quad 3! \langle \tau_{n,m} \tau_{0,1}^k \tau_{0,0}^l \rangle_3 = U(3, n+1, m, k, l | \eta_{0,1}^4) - 3U(3, n, m, k, l | \eta_{0,1}^3) + 3U(3, n-1, m, k, l | \eta_{0,1}^2).$$

3.4. There is a recursion relation for the numbers $U(g, n, m, p, q | \prod_{i=1}^t \eta_{m_i, a_i})$. Let us write it down in the most general form.

Theorem 2.

$$\begin{aligned}
(5) \quad & \left(\sum_{i=1}^t a_i \right) (2g+t-1) U(g, n, m, p, q | \prod_{i=1}^t \eta_{m_i, a_i}) = \\
& \sum_{j: m_j=0} \sum_{b_1+b_2=a_j} b_1 b_2 U(g-1, n-1, m, p, q | \eta_{1, b_1} \eta_{0, b_2} \prod_{i \neq j} \eta_{m_i, a_i}) + \\
& \sum_{j: m_j=1} \sum_{b_1+b_2=a_j} \frac{b_1 b_2}{2} U(g-1, n-1, m, p, q | \eta_{1, b_1} \eta_{1, b_2} \prod_{i \neq j} \eta_{m_i, a_i}) + \\
& \sum_{j: m_j=1} \sum_{b_1+b_2=a_j} \frac{p b_1 b_2}{6} U(g-1, n-1, m, p-1, q | \eta_{0, b_1} \eta_{0, b_2} \prod_{i \neq j} \eta_{m_i, a_i}) + \\
& \sum_{j: m_j=1} \sum_{b_1+b_2+b_3=a_j} \frac{b_1 b_2 b_3}{9} U(g-2, n-1, m, p, q | \eta_{0, b_1} \eta_{0, b_2} \eta_{0, b_3} \prod_{i \neq j} \eta_{m_i, a_i}) + \\
& \sum_{j < k: m_j=m_k=0} (a_j + a_k) U(g, n-1, m, p, q | \eta_{0, a_j+a_k} \prod_{i \neq j, k} \eta_{m_i, a_i}) + \\
& \sum_{j: m_j=0} \sum_{k: m_k=1} (a_j + a_k) U(g, n-1, m, p, q | \eta_{1, a_j+a_k} \prod_{i \neq j, k} \eta_{m_i, a_i}) + \\
& \sum_{j < k: m_j=m_k=1} \frac{p(a_j + a_k)}{3} U(g, n-1, m, p-1, q | \eta_{0, a_j+a_k} \prod_{i \neq j, k} \eta_{m_i, a_i}) + \\
& \sum_{j < k: m_j=m_k=1} \sum_{b_1+b_2=a_j+a_k} \frac{b_1 b_2}{3} U(g-1, n-1, m, p, q | \eta_{0, b_1} \eta_{0, b_2} \prod_{i \neq j, k} \eta_{m_i, a_i}) + \\
& \sum_{j < k < l: m_j=m_k=m_l=1} \frac{2(a_j + a_k + a_l)}{3} U(g, n-1, m, p, q | \eta_{0, a_j+a_k+a_l} \prod_{i \neq j, k, l} \eta_{m_i, a_i}).
\end{aligned}$$

3.5. In fact, both these theorems are very simple corollaries of the technique developed in [2]. This is discussed in details in [8]. Moreover, if we put $p = q = 0$, then both these theorems appear to be special cases of the similiar theorems in [8].

Since both these theorems could be proved just by the same argument as given in [8], we skip the proofs.

4. CALCULATIONS

4.1. In this section we prove the following theorem:

Theorem 3.

$$(6) \quad \langle \tau_{n, m} \tau_{0,1}^k \tau_{0,0}^l \rangle_3 = \frac{1}{3!12^3} \langle \tau_{n-6, m} \tau_{0,1}^{k+3} \tau_{0,0}^l \rangle_0.$$

4.2. To prove this theorem we need two lemmas.

Lemma 1.

$$(7) \quad \langle \tau_{n, m} \tau_{0,1}^k \tau_{0,0}^l \rangle_0 = \langle \tau_{n+1, m} \tau_{0,1}^k \tau_{0,0}^{l+1} \rangle_0$$

This is just the special case of the string equation (see [10]).

Lemma 2.

$$(8) \quad U(0, n, m, p, q | \prod_{i=1}^t \eta_{m_i, a_i}) = \langle \tau_{n, m} \tau_{0, 1}^p \tau_{0, 0}^q \prod_{i=1}^t \tau_{0, m_i} \rangle_0.$$

This lemma immediately follows from our definitions.

4.3. Applying many times theorem 2 and these two lemmas we obtain

$$(9) \quad U(3, n+1, m, k, l | \eta_{0, 1}^4) = \frac{317}{1548288} \langle \tau_{n-6, m} \tau_{0, 1}^{k+3} \tau_{0, 0}^l \rangle_0 + \left(\frac{11791}{23224320} + \frac{769k}{1548288} \right) \langle \tau_{n-7, m} \tau_{0, 1}^k \tau_{0, 0}^{l+1} \rangle_0;$$

$$(10) \quad U(3, n, m, k, l | \eta_{0, 1}^3) = \frac{1}{30618} \langle \tau_{n-6, m} \tau_{0, 1}^{k+3} \tau_{0, 0}^l \rangle_0 + \left(\frac{919}{7348320} + \frac{223k}{1837080} \right) \langle \tau_{n-7, m} \tau_{0, 1}^k \tau_{0, 0}^{l+1} \rangle_0;$$

$$(11) \quad U(3, n-1, m, k, l | \eta_{0, 1}^2) = \frac{k+1}{120960} \langle \tau_{n-7, m} \tau_{0, 1}^k \tau_{0, 0}^{l+1} \rangle_0.$$

4.4. It follows from these relations and theorem 1 that

$$(12) \quad 3! \langle \tau_{n, m} \tau_{0, 1}^k \tau_{0, 0}^l \rangle_3 = \frac{446}{41803776} \langle \tau_{n-6, m} \tau_{0, 1}^{k+3} \tau_{0, 0}^l \rangle_0 + \frac{19729(k+1)}{125411328} \langle \tau_{n-7, m} \tau_{0, 1}^k \tau_{0, 0}^{l+1} \rangle_0;$$

4.5. The standard calculations in genus zero (e.g., see [10]) imply the following lemma:

Lemma 3.

$$(13) \quad \langle \tau_{n, m} \tau_{0, 1}^{k+3} \tau_{0, 0}^l \rangle_0 = \frac{k+1}{3} \langle \tau_{n-1, m} \tau_{0, 1}^k \tau_{0, 0}^{l+1} \rangle_0$$

Combining equation 12 and lemma 3, we get

$$(14) \quad 3! \langle \tau_{n, m} \tau_{0, 1}^k \tau_{0, 0}^l \rangle_3 = \frac{1}{1728} \langle \tau_{n-6, m} \tau_{0, 1}^{k+3} \tau_{0, 0}^l \rangle_0.$$

This concludes the proof.

REFERENCES

- [1] E. Getzler, Topological recursion relations in genus 2, Integrable systems and algebraic geometry (Kobe/Kyoto, 1997), 73–106, World Sci. Publishing, River Edge, NJ, 1998
- [2] E.-N. Ionel, Topological recursive relations in $H^{2g}(\mathcal{M}_{g,n})$, Invent. Math. 148 (2002), no. 3, 627-658
- [3] T. J. Jarvis, T. Kimura, A. Vaintrob, Moduli spaces of higher spin curves and integrable hierarchies, Compositio Math. 126 (2001), no. 2, 157-212
- [4] M. Kontsevich, Intersection theory on the moduli space of curve and the matrix Airy function, Comm. Math. Phys. 147 (1992), 1-23
- [5] E. Looijenga, Intersection theory on Deligne-Mumford compactifications (after Witten and Kontsevich), Seminaire Bourbaki, Vol. 1992/1993, Asterisque No. 216 (1993), Exp. no. 768, 4, 187–212
- [6] A. Okounkov, R. Pandharipande, Gromov-Witten theory, Hurwitz numbers, and matrix models, Part I, arXiv: math.AG/0101147
- [7] A. Polishchuk, A. Vaintrob, Algebraic construction of Witten’s top Chern class, in Advances in Algebraic Geometry motivated by Physics, E. Previato, ed., AMS, 2001, 229-250
- [8] S. V. Shadrin, Geometry of meromorphic functions and intersections on the moduli spaces of curves, Int. Math. Res. Not. 2003, no. 38, 2051-2094
- [9] E. Witten, Two dimensional gravity and intersection theory on moduli space, Surveys in Diff. Geom. 1 (1991), 243-310
- [10] E. Witten, Algebraic geometry associated with matrix models of two dimensional gravity, Topological methods in modern mathematics (Stony Brook, NY, 1991), Publish or Perish, Houston, 1993, 235-269

STOCKHOLM UNIVERSITY AND INDEPENDENT UNIVERSITY OF MOSCOW
E-mail address: shadrin@mccme.ru